

Mott Scattering of polarized electrons in a circularly polarized laser field

B. Manaut*

Faculté polydisciplinaire, Université Sultan Moulay Slimane Béni Mellal, BP : 523, Maroc.

Y. Attaourti,[†] S. Taj, and S. Elhandi

Laboratoire de Physique des Hautes Energies et d'Astrophysique

Faculté des Sciences Semlalia, Université Cadi Ayyad, Marrakech, Boîte Postale : 2390, Maroc.

We present a study of Mott scattering of polarized electrons in the presence of a laser field with circular polarization using the helicity formalism and the introduction of the well known concept of non flip differential cross section as well as that of flip differential cross section. The results we have obtained in the presence of a laser field are coherent with those obtained in the absence of a laser field. We have studied the relativistic regime as well as the non relativistic regime that are precisely defined in the text. Two important consistency checks have been carried out successfully. The first one is that the sum of both differential cross sections (one with spin up, the other with spin down) always gives the unpolarized differential cross section. The second one is that the relativistic unpolarized differential cross section converges to the non relativistic differential cross section in the limit of small velocities. Moreover, the results obtained using the software Reduce [8] gave rise to non contracted coefficients that have been dealt with using the geometry chosen.

PACS numbers: 34.50.RK, 34.80.Qb, 12.20.Ds

I. INTRODUCTION

This work deals simultaneously with three important topics in Atomic Physics, namely the spin of a Dirac or a Dirac-Volkov particle, the concept of a spin polarized relativistic particle and finally how the laser field affects the process of scattering of such particles. We first give a brief overview of the laser. The word laser is by now well known to the lay-man because of its applications in the field of medicine (surgical and diagnostic procedures), telecommunications (fiber-optic telephone links, compact disk information storage, etc) and technology (laser drilling of materials, geodesics measurements, newspaper printing). Lasers come in different shapes, sizes and prices and under different names as ruby laser (the first laser to operate [1]), helium-neon-argon, semi-conductor laser and others. It is not possible to explain what a laser is and how it works in a few words. At this stage, it suffices to say that a laser is a light source with peculiar characteristics, drastically different from those of conventional sources such as a candle or a light bulb and is radiated in a single direction (not in all directions as in a light bulb) which enables any person working in the field to collect it in a lens and focus it in a region of very small dimensions. The spectral purity of this process and the directionality of the laser light dramatically improves the efficiency of this procedure, making it possible to concentrate a sizeable amount of power in a small region for different operations (like the melting or cutting of metals.) The laser is basically used as a very powerful light bulb. However there are others (like optical commu-

nications), in which its most important characteristics are the spectral band-width and angular aperture of the emitted beam. To understand them, one needs to consider what light is and how it is emitted, which in turn depends on the emitter, the atom. This task requires the introduction of some basic concepts of quantum mechanics. In this work, however, it is possible to describe the laser field classically since the numbers of photons involved is high [2]. As for the spin of a particle, according to Landau and Lifshitz [3], this property of elementary particles is peculiar to quantum theory and therefore has in principle no classical interpretation. In particular, it would be meaningless to imagine the "intrinsic" angular momentum of an elementary particle as being the result of its rotation about its own "axis". It is a fascinating and mysterious complication. Moreover, its practical effects prevail over the whole of Quantum Physics. The existence of spin and the statistics associated with it is the most subtle and ingenious design of Physics, without which the whole universe will collapse. Spin occupies a unique position since a wide range of Physics with different degrees of difficulty is needed for its understanding. The fact is that most textbooks (even the most advanced ones) devote at most one chapter to this subject and give only a utilitarian description. The theory of special relativity, which is essential for the understanding of spin and statistics is often forgotten except in deriving the Dirac equation [4]. The relation between spin and statistics is hard to understand. Feynman [5] wrote : " it appears to be one of the few places in Physics which can be stated very simply, but for which no one has found a simple and easy explanation. The explanation is down deep in Relativistic Quantum Mechanics [4]. This probably means that we do not have a complete understanding of the fundamental principles involved ". Now, let us turn to the physics of spin polarized free electrons. Our aim is

*manaut@fstbm.ac.ma

†attaourti@ucam.ac.ma

to oppose the current trend in science that sees specialists writing primarily for like-minded specialists. It is for example necessary to give a simple introduction of Mott Scattering one of the most important techniques in polarized electron studies. Only in recent years, has it been found possible to produce electron-beams in which the spins of the particles have a preferential direction [6]. There are many reasons for the interest in polarized electrons. One essential reason is that in physical investigations, one endeavors to define as exactly as possible the initial and/or the final states of the systems being considered and also to have electrons in the form of a well defined beam, that is a beam in which the directions of the momentum of the individual electrons are as uniform as possible. Polarization effects in electron scattering were assumed to be significant only if the electron velocity was comparable to the velocity of the light. It was not until the 1960s that large polarization effects in low-energy electron scattering were ascertained [7]. It is important, before presenting our investigation about laser-assisted Mott scattering of polarized electrons to sketch the principal steps of our treatment. For purpose of clarity and simplicity, we begin by the most basic results of Mott Scattering of polarized electrons in the absence of a laser field. Then, in the presence of a circularly polarized laser field, we give a detailed account of the formalism used and we compare the results with those obtained in the absence of a laser field. The organization of this work is as follows : in section 2, we present the scattering of polarized electrons by the Coulomb field of a heavy static charge in the absence of a laser field and we present the concept of a polarized differential cross section (in brief DCS). We also present the helicity flip DCS as well as the helicity non flip DCS. Then, we define the degree of polarization of the scattered electrons. At this stage, it is of a paramount importance to remark that in experiments, the degree of polarization of the scattered particles is measured [7]. In section 3, we present the laser assisted Mott scattering of polarized electrons in a laser field with circular polarization. After section 4, devoted to the discussion of the results obtained, we end by a brief conclusion in section 5. Throughout this work, we use atomic units $\hbar = m = e = 1$ and work with the metric tensor $g^{\mu\nu} = g_{\mu\nu} = \text{diag}(1, -1, -1, -1)$. The angular frequency we have chosen is $\omega = 0.043$ (a.u.) = 1.17 eV. It corresponds to the lasing transition of the Nd laser at a wavelength of 1064 nm.

II. MOTT SCATTERING OF POLARIZED ELECTRONS IN THE ABSENCE OF A LASER FIELD.

We begin by one of the simplest process of QED, namely the process of Mott Scattering of electrons in the lowest order of perturbation theory. We follow the usual steps of calculations and we give the transition matrix

element corresponding to this process.

$$S_{fi} = -\frac{i}{c} \int d^4x \bar{\psi}_{p_f}(x) \mathcal{A}_{coul} \psi_{p_i} \quad (1)$$

where

$$\psi_{p_i}(x) = \frac{1}{\sqrt{2E_i V}} u(p_i, s_i) e^{-ip_i x} \quad (2)$$

describes the incident electron normalized to the volume V and

$$\psi_{p_f}(x) = \frac{1}{\sqrt{2E_f V}} u(p_f, s_f) e^{-ip_f x} \quad (3)$$

describes the scattered electron with the same volume normalization. The Coulomb potential $A_{coul}^0(x)$ is generated by a static heavy nucleus of charge $-Z$, that is :

$$A_{coul}^\mu(x) = (-\frac{Z}{|\mathbf{x}|}, 0) \quad (4)$$

Therefore, we have for $\mathcal{A}_{coul}(x)$:

$$\mathcal{A}_{coul}(x) = \gamma^0 A_0 = -\frac{Z\gamma_0}{|\mathbf{x}|}$$

The transition matrix element is then given by :

$$S_{fi} = i\frac{Z}{c} \int dx^0 \int d^3\mathbf{x} \frac{\bar{u}(p_f, s_f)}{\sqrt{2E_f V}} \frac{\gamma^0}{|\mathbf{x}|} \frac{u(p_i, s_i)}{\sqrt{2E_i V}} e^{-i(p_i - p_f) \cdot x} \quad (5)$$

The integral over x_0 gives :

$$\begin{aligned} \int_{-\infty}^{+\infty} dx^0 e^{-i(p_{i0} - p_{f0}) \cdot x^0} &= \int_{-\infty}^{+\infty} d(ct) e^{-i(\frac{E_i}{c} - \frac{E_f}{c}) \cdot ct} \\ &= c \cdot 2\pi \delta(E_i - E_f) \end{aligned} \quad (6)$$

while the integral over $\mathbf{x}$ gives :

$$\int d^3x e^{i(\mathbf{p}_i - \mathbf{p}_f) \cdot \mathbf{x}} = \frac{4\pi}{(\mathbf{p}_i - \mathbf{p}_f)^2} = \frac{4\pi}{|\mathbf{q}|^2} \quad (7)$$

where we have introduced the momentum transfer $\mathbf{q} = \mathbf{p}_i - \mathbf{p}_f$. Using the standard procedures of QED [4], we get for the unpolarized DCS.

$$\frac{d\sigma}{d\Omega_f} = \frac{Z^2}{c^4} \frac{1}{|\mathbf{q}|^4} \frac{1}{2} \sum_{s_i, s_f} |\bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i)|^2 \quad (8)$$

evaluated for $E_i = E_f = E$. As a side-result this implies in turn $|\mathbf{p}_i| = |\mathbf{p}_f| = |\mathbf{p}|$. In Eq.(8) we have averaged over the initial spin polarizations s_i and summed over the final spin polarizations s_f . We obtain the simple result :

$$\frac{1}{2} \sum_{s_i, s_f} |\bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i)|^2 = 2c^2 [2\frac{E_i E_f}{c^2} - (p_i \cdot p_f) + c^2] \quad (9)$$

and with $E_i = E_f = E$ as well as $|\mathbf{p}_i| = |\mathbf{p}_f| = |\mathbf{p}|$, we end up with :

$$\frac{1}{2} \sum_{s_i, s_f} |\bar{u}(\mathbf{p}_f, s_f) \gamma^0 u(\mathbf{p}_i, s_i)|^2 = 2c^2 \left[\frac{E^2}{c^2} + |\mathbf{p}|^2 \cos(\theta) + c^2 \right] \quad (10)$$

where θ is the scattering angle and $(p_i \cdot p_f) = E^2/c^2 - \mathbf{p}_f \cdot \mathbf{p}_i = E^2/c^2 - |\mathbf{p}|^2 \cos(\theta)$. Using the relation $\beta^2 E^2 = c^2 |\mathbf{p}|^2$ ($\beta = \frac{v}{c}$) and $|\mathbf{q}| = |\mathbf{p}_i - \mathbf{p}_f| = 2|\mathbf{p}| \sin(\theta/2)$, the final expression for the unpolarized DCS is :

$$\begin{aligned} \frac{d\bar{\sigma}}{d\Omega_f}_{Mott} &= \frac{1}{4} \frac{Z^2}{c^2 \beta^2 |\mathbf{p}|^2} \frac{1 - \beta^2 \sin^2(\theta/2)}{\sin^4(\theta/2)} \\ &= \frac{d\bar{\sigma}}{d\Omega_f}_{Ruth} (1 - \beta^2 \sin^2(\theta/2)) \end{aligned} \quad (11)$$

where

$$\frac{d\bar{\sigma}}{d\Omega_f}_{Ruth} = \frac{Z^2}{4c^2 \beta^2 |\mathbf{p}|^2} \frac{1}{\sin^4(\theta/2)} \quad (12)$$

is the Rutherford unpolarized DCS in the limit of small velocities ($\beta \rightarrow 0$). The calculations carried out up to now assumed that the initial electron spin is not observed. So, we focus now on the scattering of polarized Dirac particles. For that purpose, we need some formalism to describe the spin polarization [4]. Free electrons with 4-momentum p and spin s are described by the free spinors $u(p, s)$. The spin 4-vector s^μ is defined by :

$$s^\mu = \frac{1}{c} (|\mathbf{p}|, \frac{E}{c} \hat{p}) \quad (13)$$

where $\hat{p} = \mathbf{p}/|\mathbf{p}|$ is the unit vector defining the direction of the 3-vector $\mathbf{p}$. The 4-vector s^μ is a Lorentz vector in a frame in which the particle moves with momentum $\mathbf{p}$. The following properties can be easily checked :

$$s^\mu \cdot s_\mu = -1 \quad (14)$$

This is the normalization relation for the 4-vector s^μ . One also have :

$$(p \cdot s) = p^\mu \cdot s_\mu = 0 \quad (15)$$

It is the orthogonality relation between p and s . Next, we introduce the spin projection operator :

$$\hat{\Sigma}(s) = \frac{1}{2} (1 + \gamma_5 \not{s}) \quad (16)$$

with $\gamma_5 = i\gamma^0 \gamma^1 \gamma^2 \gamma^3 = -i\gamma_0 \gamma_1 \gamma_2 \gamma_3$. This operator has a simple property :

$$\hat{\Sigma}(s) u(p, +s) = u(p, +s) \quad , \quad \hat{\Sigma}(s) u(p, -s) = 0 \quad (17)$$

This formalism can be applied to helicity states where the direction of the spin points along the direction of the momentum 3-vector $\mathbf{p}$.

$$S'_\lambda = \lambda \frac{\mathbf{p}}{|\mathbf{p}|} \quad \lambda = \pm 1 \quad (18)$$

Therefore, the definition of a 4-spin vector follows :

$$S^\mu_\lambda = \frac{\lambda}{c} (|\mathbf{p}|, \frac{E}{c} \hat{p}) \quad (19)$$

We calculate now the polarized DCS for Mott Scattering of an electron with a well defined momentum p_i and a well defined spin s_i . If also the final spin s_f is measured, the polarized DCS reads :

$$\frac{d\sigma}{d\Omega_f} = \frac{Z^2}{c^2} \frac{|\bar{u}(p_f, s_f) \gamma^0 u(p_i, s_i)|^2}{|\mathbf{q}|^4} \quad (20)$$

In Eq.(20), one has to be careful since there are no summation on spin (either initial or final) polarizations.

We introduce the two operators :

$$\hat{\Sigma}_{\lambda_i}(s_i) = \frac{1}{2} (1 + \lambda_i \gamma_5 \not{s}_i) \quad (21)$$

$$\hat{\Sigma}_{\lambda_f}(s_f) = \frac{1}{2} (1 + \lambda_f \gamma_5 \not{s}_f) \quad (22)$$

and we obtain for the polarized DCS :

$$\begin{aligned} \frac{d\sigma}{d\Omega_f}(\lambda_i, \lambda_f) &= \frac{Z^2}{c^4} \frac{1}{4} Tr \{ \gamma^0 (1 + \lambda_i \gamma_5 \not{s}_i) (\not{p}_i c + c^2) \\ &\quad \gamma^0 (1 + \lambda_f \gamma_5 \not{s}_f) (\not{p}_f c + c^2) \} \end{aligned} \quad (23)$$

using the relations $(p_i \cdot s_f) = \frac{E}{c^2} |\mathbf{p}| (1 - \cos(\theta)) = (p_f \cdot s_i)$, $(s_i \cdot s_f) = (|\mathbf{p}|^2 - E^2 \cos(\theta)/c^2)/c^2$ as well as $(p_i \cdot p_f) = \frac{E^2}{c^2} - |\mathbf{p}|^2 \cos(\theta)$ we obtain :

$$\begin{aligned} \frac{d\sigma}{d\Omega_f}(\lambda_i, \lambda_f) &= \frac{Z^2}{|\mathbf{q}|^4 c^4} 2 \left\{ E^2 \cos^2(\theta/2) + c^4 \sin^2(\theta/2) \right. \\ &\quad \left. + \lambda_i \lambda_f (E^2 \cos^2(\theta/2) - c^4 \sin^2(\theta/2)) \right\} \end{aligned} \quad (24)$$

At this stage, let us note that the λ_i and λ_f have the following properties :

$$\lambda_i^2 = \lambda_f^2 = 1 \quad (25)$$

$$\lambda_i \cdot \lambda_f = 1 \implies \begin{cases} \lambda_i = \lambda_f = 1 \\ \lambda_i = \lambda_f = -1 \end{cases} \quad (26)$$

$$\lambda_i \cdot \lambda_f = -1 \implies \begin{cases} \lambda_i = -\lambda_f = 1 \\ \lambda_i = -\lambda_f = -1 \end{cases} \quad (27)$$

So, if during the scattering process, $\lambda_f = -\lambda_i$ which means that a helicity flip occurred, the flip polarized DCS is :

$$\begin{aligned} \frac{d\sigma}{d\Omega_f} \Big|_{flip} &= \frac{4Z^2}{|\mathbf{q}|^4 c^4} c^4 \sin^2(\theta/2) \\ &= \frac{d\sigma}{d\Omega_f} \Big|_{Ruth} \frac{c^4}{E^2} \sin^2(\theta/2) \end{aligned} \quad (28)$$

The case where there is no helicity flip corresponds to $\lambda_i = \lambda_f$, so that $\lambda_i \lambda_f = 1$ and the helicity non flip polarized DCS is given by :

$$\left. \frac{d\sigma}{d\Omega_f} \right|_{nonflip} = \left. \frac{d\sigma}{d\Omega_f} \right|_{Ruth} \cos^2(\theta/2) \quad (29)$$

Of course, the sum of the helicity flip polarized DCS and the helicity non flip polarized DCS must give the unpolarized DCS.

$$\left. \frac{d\bar{\sigma}}{d\Omega_f} \right|_{Mott} = \left. \frac{d\sigma}{d\Omega_f} \right|_{nonflip} + \left. \frac{d\sigma}{d\Omega_f} \right|_{flip} \quad (30)$$

Noting that $E^2 = |\mathbf{p}|^2 c^2 + c^4$ or $c^4 = E^2 - |\mathbf{p}|^2 c^2$ we have :

$$\frac{c^4}{E^2} \sin^2(\theta/2) + \cos^2(\theta/2) = 1 - \beta^2 \sin^2(\theta/2) \quad (31)$$

and the final results is :

$$\left. \frac{d\bar{\sigma}}{d\Omega_f} \right|_{Mott} = \left. \frac{d\sigma}{d\Omega_f} \right|_{Ruth} [1 - \beta^2 \sin^2(\theta/2)] \quad (32)$$

This result is of a paramount importance.

For every process, the sum of the flip polarized DCS and the non flip polarized DCS always gives the unpolarized DCS. In fact, this is theoretically exact but, as we shall see very soon (section 3), when it comes to numerical simulations and where infinite summations over the number of laser photons are involved, due to a unavoidable lack of precision, the theoretical exact result becomes a numerical approximate result. Another important result is the degree of polarization defined by :

$$P = \frac{\left. \frac{d\sigma}{d\Omega_f} \right|_{\lambda_i = \lambda_f = 1} - \left. \frac{d\sigma}{d\Omega_f} \right|_{\lambda_i = -\lambda_f = 1}}{\left. \frac{d\sigma}{d\Omega_f} \right|_{\lambda_i = \lambda_f = 1} + \left. \frac{d\sigma}{d\Omega_f} \right|_{\lambda_i = -\lambda_f = 1}} \quad (33)$$

For our process, P reads :

$$\begin{aligned} P &= \frac{\cos^2(\theta/2) - \frac{c^4}{E^2} \sin^2(\theta/2)}{\cos^2(\theta/2) + \frac{c^4}{E^2} \sin^2(\theta/2)} \\ &= 1 - 2 \frac{c^4 \sin^2(\theta/2)}{E^2 \cos^2(\theta/2) + c^4 \sin^2(\theta/2)} \end{aligned} \quad (34)$$

and with $E = \gamma c^2$, one has :

$$P = 1 - 2 \frac{\sin^2(\theta/2)}{\gamma^2 \cos^2(\theta/2) + \sin^2(\theta/2)} \quad (35)$$

We notice that for the process of Mott scattering in the absence of the laser field, only two parameters are relevant : the relativistic parameter $\gamma = 1/\sqrt{1 - \beta^2}$ and the scattering angle θ_{if} . Moreover, one can see from Fig. (1) that when the collision approaches the relativistic domain, the degree of polarization becomes less strongly dependent on the relativistic parameter γ and becomes




FIG. 1: Degree of polarization as function of the scattering angle θ_{if} , for different values of the relativistic parameter γ .

close to a constant value $P \simeq 1$. The value $P = 1$ corresponds to a collision when no spin flip occurs whereas the value $P = -1$ corresponds to the reverse situation (there is a spin flip). In Fig. (2), we plot P as a function of the final electron momentum θ_f and a different picture emerges. Since the scattering angle is not involved, the minimum value of P shifts to a value greater than $P = -1$ and this trend is more and more noticeable for increasing values of γ .

III. MOTT SCATTERING OF POLARIZED ELECTRONS IN THE PRESENCE OF A CIRCULARLY POLARIZED LASER FIELD.

The 4-vector potential A^μ chosen is such that :

$$A^\mu = a_1^\mu \cos(\phi) + a_2^\mu \sin(\phi) \quad (36)$$

with $(a_1.a_1) = -|\mathbf{a}|^2 = -|\mathbf{A}|^2$, $(a_2.a_2) = -|\mathbf{a}|^2 = -|\mathbf{A}|^2$ and the polarization 4-vectors a_1^μ and a_2^μ are such that :

$$a_1^\mu = |\mathbf{a}| e_x^\mu = |\mathbf{a}| (0, 1, 0, 0)$$

and

$$a_2^\mu = |\mathbf{a}| e_y^\mu = |\mathbf{a}| (0, 0, 1, 0)$$




FIG. 2: Degree of polarization as function of the angle θ_f , for different values of the relativistic parameter γ .

In Eq.(36), $\phi = (k.x) = k_\mu.x^\mu$ with $k^\mu = (k^0, 0, 0, k^3) = \omega/c(1, 0, 0, 1)$. A^μ satisfies the Lorentz condition :

$$k_\mu.A^\mu = 0 \quad (37)$$

which implies $(k.a_1) = (k.a_2) = 0$ i.e, we choose $\mathbf{k}$ along the direction of the Oz axes.

Now, in presence of a laser field, the Dirac wave functions describing the incident and scattered electrons have to be replaced by the Dirac-Volkov wave function [3] which are given by :

$$\psi_q(x) = R(q) \frac{u(p, s)}{\sqrt{2QV}} e^{iS(q, s)} \quad (38)$$

where

$$R(q) = R(p) = 1 + \frac{k.A}{2c(k.q)} = 1 + \frac{k.A}{2c(k.p)} \quad (39)$$

since the 4-vector q^μ that now replaces p^μ is such that :

$$q^\mu = p^\mu - \frac{\bar{A}^2}{2c^2(k.p)} k^\mu \quad (40)$$

The time component of q^μ is such that :

$$q^0 = \frac{Q}{c} = \frac{E}{c} - \frac{a^2}{2c^2(k.p)} \frac{\omega}{c} = \frac{1}{c} (E - \frac{a^2 \omega}{2c^2(k.p)}) \quad (41)$$

We shall see that this 4-vector is the new canonical impulsion 4-vector that the electron feels inside the laser field. The 4-vector q^μ is called the quasi-impulsion and has the following property :

$$\begin{aligned} q_\mu.q^\mu &= [p - \frac{a^2 k}{2c^2(k.p)}]_\mu [p - \frac{a^2 k}{2c^2(k.p)}]^\mu \\ &= p_\mu.p^\mu - \frac{a^2}{c^2} \\ &= c^2(1 - \frac{a^2}{c^4}) \end{aligned} \quad (42)$$

Remark also that we can write :

$$R(q) = [1 + c(q)k.A] = [1 + c(p)k.A] \quad (43)$$

with $c(q) = c(p) = 1/[2c(k.q)] = 1/[2c(k.p)]$, since $(k.q) = (k.p)$ and $k^2 = k^\mu.k_\mu = 0$. The function $S(q, s)$ appearing in Eq.(38) is such that :

$$S(q, s) = -q.x - i \frac{(p.a_1)}{c(k.p)} \sin(\phi) + i \frac{(p.a_2)}{c(k.p)} \cos(\phi) \quad (44)$$

One has for the transition matrix element :

$$S_{fi} = -\frac{i}{c} \int d^4x \bar{\psi}_{q_f}(x) A_{coul}(x) \psi_{q_i}(x) \quad (45)$$

One must be very cautious not to confuse the Coulomb 4-vector potential due to the charge of the nucleus with the 4-vector potential of the circularly polarized laser field. As before one has $A_{coul}(x) = -Z\gamma^0/|\mathbf{x}|$ and therefore :

$$S_{fi} = i \frac{Z}{c} \int d^4x \bar{\psi}_{q_f}(x) \frac{\gamma^0}{|\mathbf{x}|} \psi_{q_i}(x) \quad (46)$$

with

$$\psi_{q_i}(x) = R(p_i) \frac{u(p_i, s_i)}{\sqrt{2QV}} e^{iS(q_i, x)} \quad (47)$$

and

$$\psi_{q_f}(x) = \frac{\bar{u}(p_f, s_f)}{\sqrt{2QV}} \bar{R}(p_f) e^{-iS(q_f, x)} \quad (48)$$

we transform $e^{iS(q_i, x) - iS(q_f, x)}$, if we introduce :

$$z = \sqrt{\alpha_1^2 + \alpha_2^2} \quad (49)$$

with

$$\alpha_1 = \frac{1}{c} \left(\frac{(a_1.p_i)}{(k.p_i)} - \frac{(a_1.p_f)}{(k.p_f)} \right) \quad (50)$$

and

$$\alpha_2 = \frac{1}{c} \left(\frac{(a_2.p_i)}{(k.p_i)} - \frac{(a_2.p_f)}{(k.p_f)} \right) \quad (51)$$

we get :

$$S(q_i, x) - S(q_f, x) = -(q_i - q_f).x - z \sin(\phi - \phi_0) \quad (52)$$

where :

$$\cos(\phi_0) = \frac{\alpha_1}{z} \quad \text{or} \quad \sin(\phi_0) = \frac{\alpha_2}{z} \quad (53)$$

Now, let us look closely at the quantities in Eq.(46). First, we have :

$$\bar{\psi}_{q_f}(x)\gamma^0\psi_{q_i}(x) = \bar{u}(p_f, s_f)\bar{R}(p_f)\gamma^0 R(p_i)u(p_i, s_i) \quad (54)$$

In particular, we remark that :

$$\begin{aligned} \bar{R}(p_f)\gamma^0 R(p_i) &= \{1 + c(p_f)[\phi_1 k \cos \phi + \phi_2 k \sin \phi]\} \\ &\quad \times \gamma^0 \{1 + c(p_i)[k \phi_1 \cos \phi + k \phi_2 \sin \phi]\} \\ &= \{\gamma^0 + c(p_f)[\phi_1 k \gamma^0 \cos \phi + \phi_2 k \gamma^0 \sin \phi]\} \\ &\quad \times \{1 + c(p_i)[k \phi_1 \cos \phi + k \phi_2 \sin \phi]\} \quad (55) \end{aligned}$$

Before invoking the well known relations involving ordinary Bessel functions, we have first to transform Eq.(55) in a more compact form. Eq.(55) contains nine terms, and after some algebraic calculations the final result for $\bar{R}(q_f)\gamma^0 R(q_i) = \bar{R}(p_f)\gamma^0 R(p_i)$ is :

$$\begin{aligned} \bar{R}(q_f)\gamma^0 R(q_i) &= \gamma^0 - 2\frac{\omega}{c}c(p_i)c(p_f)a^2k \\ &\quad + \{c(p_i)\gamma^0 k \phi_1 + c(p_f)\phi_1 k \gamma^0\} \cos \phi \\ &\quad + \{c(p_i)\gamma^0 k \phi_2 + c(p_f)\phi_2 k \gamma^0\} \sin \phi \\ &= C_0 + C_1 \cos(\phi) + C_2 \sin(\phi) \quad (56) \end{aligned}$$

with

$$\begin{aligned} C_0 &= \gamma^0 - 2\frac{\omega}{c}c(p_i)c(p_f)a^2k \\ C_1 &= c(p_i)\gamma^0 k \phi_1 + c(p_f)\phi_1 k \gamma^0 \\ C_2 &= c(p_i)\gamma^0 k \phi_2 + c(p_f)\phi_2 k \gamma^0 \end{aligned}$$

Now, we have :

$$\left\{ \begin{array}{c} 1 \\ \cos(\phi) \\ \sin(\phi) \end{array} \right\} e^{-iz \sin(\phi - \phi_0)} = \sum_{s=-\infty}^{\infty} \left\{ \begin{array}{c} B_n(z) \\ B_{1n}(z) \\ B_{2n}(z) \end{array} \right\} e^{-in\phi} \quad (57)$$

with

$$\begin{cases} B_n(z) = J_n(z)e^{in\phi_0} \\ B_{1n}(z) = (J_{n+1}(z)e^{i(n+1)\phi_0} + J_{n-1}(z)e^{i(n-1)\phi_0})/2 \\ B_{2n}(z) = (J_{n+1}(z)e^{i(n+1)\phi_0} - J_{n-1}(z)e^{i(n-1)\phi_0})/2i \end{cases} \quad (58)$$

Therefore :

$$\begin{aligned} \bar{R}(p_f)\gamma^0 R(p_i) &= \sum_{n=-\infty}^{+\infty} [C_0 B_n(z) + C_2 B_{1n}(z) \\ &\quad + C_3 B_{2n}(z)] e^{-in\phi} \\ &= \sum_{n=-\infty}^{+\infty} \Gamma_n e^{-in\phi} \quad (59) \end{aligned}$$

The transition matrix element becomes :

$$S_{fi} = iZ2\pi\delta(Q_i - Q_f + n\omega) \frac{4\pi}{|\mathbf{q}|^2} \frac{\bar{u}(p_f, s_f)}{\sqrt{2Q_f V}} \Gamma_n \frac{u(p_i, s_i)}{\sqrt{2Q_i V}} \quad (60)$$

Using the standard procedures of QED, one obtains for the polarized DCS, we have :

$$\frac{d\sigma}{d\Omega_f} = \sum_{n=-\infty}^{+\infty} \frac{d\sigma^{(n)}}{d\Omega_f} \quad (61)$$

where :

$$\frac{d\sigma^{(n)}}{d\Omega} = \frac{|q_f|}{|q_i|} \frac{Z^2}{c^4} \frac{1}{|\mathbf{q}|^4} |\bar{u}(p_f, s_f)\Gamma_n u(p_i, s_i)|^2 \quad (62)$$

evaluated for $Q_f = Q_i + n\omega$.
The squared matrix element is :

$$\begin{aligned} |\bar{u}(p_f, s_f)\Gamma_n u(p_i, s_i)|^2 &= u^\dagger(p_i, s_i)\Gamma_n^\dagger \gamma^0 u(p_f, s_f) \bar{u}(p_f, s_f)\Gamma_n u(p_i, s_i) \\ &= \bar{u}(p_i, s_i)\gamma^0 \Gamma_n^\dagger \gamma^0 u(p_f, s_f) \bar{u}(p_f, s_f)\Gamma_n u(p_i, s_i) \quad (63) \end{aligned}$$

If the operators $\hat{\Sigma}_{\lambda_i}(s_i)$ and $\hat{\Sigma}_{\lambda_f}(s_f)$ are introduced, we obtain :

$$|\bar{u}(p_f, s_f)\Gamma u(p_i, s_i)|^2 = Tr \left\{ \Gamma_n \frac{(1 + \lambda_i \gamma_5 \not{p}_i)}{2} (\not{p}_i c + c^2) \bar{\Gamma}_n \frac{(1 + \lambda_f \gamma_5 \not{p}_f)}{2} (\not{p}_f c + c^2) \right\} \quad (64)$$

where :

$$\begin{aligned} \bar{\Gamma}_n &= \gamma^0 \Gamma_n^\dagger \gamma^0 \\ &= [\gamma^0 - 2\frac{\omega}{c}c(p_f)c(p_i)a^2k] B_n^* \\ &\quad + [c(p_i)\phi_1 k \gamma^0 + c(p_f)\gamma^0 k \phi_1] B_{1n}^* \\ &\quad + [c(p_i)\phi_2 k \gamma^0 + c(p_f)\gamma^0 k \phi_2] B_{2n}^* \quad (65) \end{aligned}$$

IV. RESULTS AND DISCUSSIONS.

Before presenting the results and their physical interpretation, we would like to emphasize the fact that the RE-

DUCE code we have written gave very long analytical expressions which were difficult to incorporate in the corresponding FORTRAN code we wrote to extract figures and tables. From now on, the pair of indices n, f will stand for "non flip" and the index f alone will stand for "flip". For the non flip DCS, let us simply write

$$\left. \frac{d\sigma}{d\Omega} \right|_{n,f} = \sum_{n=-\infty}^{n=\infty} \left. \frac{d\sigma^{(n)}}{d\Omega} \right|_{n,f} \quad (66)$$

where

$$\begin{aligned} \left. \frac{d\sigma^{(n)}}{d\Omega_f} \right|_{(n,f)} &= \frac{|\mathbf{q}_f|}{|\mathbf{q}_i|} \frac{Z^2}{c^4} \frac{1}{|\mathbf{q}|^4} \left\{ \mathcal{A}(\lambda_i = \lambda_f = 1) J_n^2(z) \right. \\ &\quad + \mathcal{B}(\lambda_i = \lambda_f = 1) [J_{n+1}^2(z) + J_{n-1}^2(z)] \\ &\quad + \mathcal{C}(\lambda_i = \lambda_f = 1) J_{n+1}(z) J_{n-1}(z) \\ &\quad \left. + \mathcal{D}(\lambda_i = \lambda_f = 1) J_n(z) [J_{n+1}(z) + J_{n-1}(z)] \right\} \end{aligned} \quad (67)$$

The four coefficients $\mathcal{A}(\lambda_i, \lambda_f)$, $\mathcal{B}(\lambda_i, \lambda_f)$, $\mathcal{C}(\lambda_i, \lambda_f)$ and $\mathcal{D}(\lambda_i, \lambda_f)$ are very long to write down so we prefer to focus on their global contents instead of giving tedious details and explanations. First, we noticed that in both coefficients $\mathcal{A}_{n,f} = \mathcal{A}(\lambda_i = \lambda_f = 1)$ and $\mathcal{C}_{n,f} = \mathcal{C}(\lambda_i = \lambda_f = 1)$, there is no occurrence of the various completely anti-symmetric tensors $\varepsilon_{\alpha\beta\gamma\delta}$ (where the indices are Lorentz ones and take integer values from 0 to 3). This clearly means that these tensors were totally contracted and we ended up with two tractable coefficients that were very easy to incorporate into the main FORTRAN code. Let us remind the reader that we used throughout this work, the convention :

$$\varepsilon_{0123} = 1 \quad (68)$$

meaning that $\varepsilon_{\alpha\beta\gamma\delta} = 1$ for an even permutation of the Lorentz indices whereas $\varepsilon_{\alpha\beta\gamma\delta} = -1$ for an odd permutation of the Lorentz indices and finally $\varepsilon_{\alpha\beta\gamma\delta} = 0$ otherwise. Second, the coefficients $\mathcal{B}_{n,f} = \mathcal{B}(\lambda_i = \lambda_f = 1)$ as well as $\mathcal{D}_{n,f} = \mathcal{D}(\lambda_i = \lambda_f = 1)$ contained various non contracted tensors. For example in $\mathcal{B}_{n,f}$, there are thirty one non contracted tensors involving $\varepsilon_{\alpha\beta\gamma\delta}$ whereas in $\mathcal{D}_{n,f}$, there are sixty four. Particle physicists are very often dealing with these. As for us, we were confronted for the first time with coefficients like for example in $\mathcal{B}_{n,f}$:

$$\varepsilon(a_1, a_2, k, p_i) = \varepsilon_{\alpha\beta\gamma\delta} a_1^\alpha a_2^\beta k^\gamma p_i^\delta \quad (69)$$

and

$$\varepsilon(a_1, a_2, k, v) = \varepsilon_{\alpha\beta\gamma\delta} a_1^\alpha a_2^\beta k^\gamma v^\delta \quad (70)$$

Such coefficients are, at first sight, very complicated to evaluate. But, when following the conventions of A.G Grozin [8], we found that the only non-vanishing contribution of a_1 corresponds to the Lorentz index $\alpha = 1$ while the only non-vanishing contribution of a_2 corresponds to




FIG. 3: Envelope of the relativistic DCS scaled in 10^{-14} a.u. as a function of the energy transfer scaled in units of the laser photon energy ω for an electrical field strength of $\mathcal{E} = 1$ a.u. and a relativistic parameter $\gamma = 2$.

the Lorentz index $\beta = 2$. These helpful values are due to the choice of the 4-potential A^μ . Also, it allows to deduce that the only non trivial indices remaining for k^γ and p_i^δ are the pairs $(\gamma = 0, \delta = 3)$ and $(\gamma = 3, \delta = 0)$. So, the non contracted tensor $\varepsilon(a_1, a_2, k, p_i)$ reduces to :

$$\begin{aligned} \varepsilon(a_1, a_2, k, p_i) &= |\mathbf{a}|^2 [\varepsilon_{1203} k^0 \mathbf{p}_i^3 + \varepsilon_{1230} k^3 p_i^0] \\ &= |\mathbf{a}|^2 \frac{\omega}{c} \left[|\mathbf{p}_i| \cos \theta_i - \frac{E_i}{c} \right] \end{aligned} \quad (71)$$

where θ_i is the angle between the initial electron momentum and the Oz axis. The coefficient $\varepsilon(a_1, a_2, k, v)$ is easier to evaluate since the Lorentz index corresponding to $v = (1, 0, 0, 0)$ is zero $\delta = 0$. This trick is often used in all QED textbooks [4]. So, the only choice left for k is $\gamma = 3$. Thus, we have :

$$\begin{aligned} \varepsilon(a_1, a_2, k, v) &= \varepsilon_{1230} a_1^1 a_2^2 k^3 v^0 \\ &= -|\mathbf{a}|^2 \frac{\omega}{c} \end{aligned} \quad (72)$$

It is not necessary to give the whole set of the coefficients involving the non contracted tensors appearing in $\mathcal{B}_{n,f}$ and $\mathcal{D}_{n,f}$. It is sufficient to follow the rules concerning the tensor $\varepsilon_{\alpha\beta\gamma\delta}$ and the geometry chosen for $\mathbf{a}_1$, $\mathbf{a}_2$ and $\mathbf{k}$, bearing in mind that $a_1^\mu = |\mathbf{a}| e_1^\mu = |\mathbf{a}|(0, 1, 0, 0)$, $a_2^\mu = |\mathbf{a}| e_2^\mu = |\mathbf{a}|(0, 0, 1, 0)$ and finally $k^\sigma = \frac{\omega}{c}(1, 0, 0, 1)$.

The same holds for the coefficients $\mathcal{B}_f = \mathcal{B}(\lambda_i = -\lambda_f = 1)$ and $\mathcal{D}_f = \mathcal{D}(\lambda_i = -\lambda_f = 1)$. The complete expressions of the coefficients $\mathcal{A}$, $\mathcal{B}$, $\mathcal{C}$ and $\mathcal{D}$ in both cases (nf, f) can explicitly be found in the Appendix. It is important to remind that these coefficients occur only in presence of a laser field. If we put $A^\mu = (0, \mathbf{0})$, we easily recover the results of section 2. Now, let us turn to the discussion of the results concerning $d\sigma/d\Omega_f|_{n,f}$ and $d\sigma/d\Omega_f|_f$. We chose two regimes : a) the relativistic regime corresponding to a relativistic parameter $\gamma = 2$ and an electric field strength $\mathcal{E} = 1$ a.u and b) the non relativistic regime corresponding to a relativistic parameter $\gamma = 1.0053$ and an electric field strength $\mathcal{E} = 0.05$ a.u. There are two consistency checks that must be made. First, we have to show numerically that the sum of $(DCS)_f$ and $(DCS)_{n,f}$ always gives the unpolarized DCS and second, the non relativistic description for the unpolarized DCS must give the unpolarized DCS we have found in the formalism we have developed.

A. The relativistic regime




FIG. 4: Degree of polarization P for an electrical field strength of $\mathcal{E} = 1$ a.u and a relativistic parameter $\gamma = 2$.

This regime corresponds to a total energy of the incoming electron $E_i = 0.510858$ MeV which is the rest energy of the electron so, it is located below the threshold energy needed for the process of pair creation. As for the cutoffs,

the negative part of the spectrum corresponds approximately to -150000 photons (emitted by the laser). As a side-result, the various DCS (n, f or f) decrease abruptly because the arguments of the Bessel functions are close to their indices. Once again, there is an asymmetry between the emission part of the envelope and the absorption part of the same envelope for the unpolarized DCSs because of the presence of the denominator $|\mathbf{q}|^4$ in Eq.(67). Now, the positive part of the spectrum corresponds approximately to 175000 photons absorbed by the laser field. In Fig. (3), we show the envelope of $d\sigma/d\Omega_f$ as a function of the net number of photons exchanged. The cutoffs are $n \simeq -150000$ photons for the negative part of the envelope and $n_{max} \simeq 170000$ photons for the positive part of the envelope that corresponds to the absorptive part of the spectrum. The geometry chosen for Fig. (3) is $\theta_i = 45^\circ$, $\phi_i = 0^\circ$, $\theta_f = 50^\circ$ and $\phi_f = 90^\circ$. One may ask legitimately if such cutoffs are geometry-dependent. The answer is that indeed they are geometry dependent since for a different choice of the initial, final momentum angular coordinates $\mathbf{p}_i$ and $\mathbf{p}_f$ namely $\theta_i = 30^\circ$, $\phi_i = 0^\circ$, $\theta_i = 75^\circ$ and $\phi_f = 90^\circ$ the cutoffs obtained are $n_{min} \simeq -147000$ photons and $n_{max} \simeq +154000$ photons. In Fig. (4), we show the degree of polarization P corresponding to this regime.




FIG. 5: The various relativistic DCSs scaled in 10^{-11} a.u as a function of the angle θ_f in degrees for an electrical field strength of $\mathcal{E} = 1$ a.u and a relativistic parameter $\gamma = 2$. The corresponding number of photons exchanged is ± 500 .




FIG. 6: First consistency check related to the sum of DCS(up) and DCS(down) compared to the unpolarized DCS scaled in 10^{-11} a.u. as a function of the angle θ_f in degrees for an electrical field strength of $\mathcal{E} = 1$ a.u. and a relativistic parameter $\gamma = 2$. The corresponding number of photons exchanged is ± 500 .

For this degree of polarization, we obtained a qualitative result similar to that of a linearly polarized field. We reached the same conclusion as in [9] that the degree of polarization.

$$P = 1 - \frac{d\sigma/d\Omega_f(\downarrow)}{[d\sigma/d\Omega_f(\uparrow) + d\sigma/d\Omega_f(\downarrow)]} \quad (73)$$

is weakly dependent on the number of photons exchanged. This degree of polarization P varies as a function of the angle θ_f for the following geometry ($\theta_i = 45^\circ$, $\phi_i = 0$, $\phi_f = 90^\circ$ and $0 \leq \theta_f \leq 180^\circ$). To begin with, we have made simulations concerning the various DCSs for a set of net number of photons exchanged. These sets (± 100 , ± 200 , ± 300 , ± 400 , ± 500 , ± 1000) showed that the order of magnitude of the non flip DCS is close to the unpolarized DCS but the contribution of the flip DCS is not completely negligible even if it is small compared to both. The behavior of the three DCSs when the number of photons exchanged increases has an influence over the numerical values of the differential cross sections but as we are limited in our computational capabilities, it is not possible to achieve numerical convergence but the most important result is that the sum of DCS(up) and DCS(down) always gives the unpolarized DCS which is

a very important consistency check. Both Figs. (5,6) are accurate illustrations of this consistency check.

B. The non relativistic regime.

In this regime, the dressing of the angular coordinates of $\mathbf{p}_i$ as well as those of $\mathbf{p}_f$ is not important. The second and final consistency check is shown in Fig. (7). First, we give the non relativistic DCS obtained by using Schrödinger-Volkov wave functions :

$$\frac{d\sigma^{(n)}}{d\Omega_f} = \frac{4Z^2|\mathbf{p}_f|}{|\mathbf{p}_i|} J_n^2\left(\frac{a}{c\omega} \sqrt{\Delta.\hat{x} + \Delta.\hat{y}}\right) \frac{1}{|\mathbf{p}_i - \mathbf{p}_f + n\mathbf{k}|^4} \quad (74)$$

The two DCSs are very close, which was to be expected but with small deviations coming from the unpolarized relativistic DCS in the limit of small velocities and this is due to the fact that error propagation is more likely to occur when one is dealing with very long expressions. This is indeed the case for the latter. We have summed over ± 500 photons to obtain the curves in Fig. (7). There is a




FIG. 7: Second consistency check in the non relativistic regime related to the sum of DCS(up) and DCS(down) compared to the non relativistic DCS scaled in 10^{-4} a.u. as a function of the angle θ_f in degrees for an electrical field strength of $\mathcal{E} = 0.05$ a.u. and a relativistic parameter $\gamma = 1.0053$. The corresponding number of photons exchanged is ± 500 . For large negative and positive angles, it becomes scaled in 10^{-5} a.u.

very good argument between the two curves and it is possible by increasing the number of net photons exchanged to reach the same value as that of the unpolarized DCS in absence of the laser field. Many similar results were obtained because of a pseudo sum-rule that was shown by Bunkin and Fedorov as well as Kroll and Watson [10].




FIG. 8: Envelope of the two unpolarized DCS in the non relativistic regime for a number of net photons exchanged ± 500 scaled in units of 10^{-7} . The two curves are indistinguishable and are shifted from the elastic peak corresponding to $\theta_f = 0$. The cutoffs are visible.

Even if the non relativistic description does not take into account the spin of the electron, the agreement between the two DCSs shows in the one hand, that spin effects are not important in this regime while on the other hand, given the more complete and sophisticated relativistic description, the non relativistic limit of the relativistic DCS always gives results very close to the non

relativistic description. This does not mean that spin effects are irrelevant but only that their contribution to the relativistic DCS (for small velocities) is small enough to be noticeable. As for the cutoffs, one can observe from Fig. (8) that for the negative part of the spectrum, it is located approximately at -450 photons while for the absorptive part of the spectrum, it is located approximately at $+470$ photons.

V. CONCLUSION.

In this work, we have checked that results already known in the absence of a laser field are also valid in presence of a laser field, at least in first order of perturbation theory. The motivation of this study was to compare the order of magnitude for the various DCSs in the relativistic regime that we already found in [9] particularly in the case of a laser field with linear polarization and we have found (see Fig. (6)) that indeed when scaled in $10^{(-11)}$, both DCSs are of similar order of magnitude and shape. Moreover, when focusing only on the formalism for polarized electrons, the main difference between the linear and the circular polarization of the laser field, is that a thorough analysis of the contents of the formal expression of DCS(up) and DCS(down) is more complicated in the case of the circular polarization than the linear polarization of the laser field and this is due to the fact that these two DCSs contain non contracted symbols that we had to deal with for the first time. We succeeded in the simulations of the main two consistency checks namely that the sum of DCS(up) and DCS(down) always gives the unpolarized DCS regardless of the number of net photons exchanged and also that in the non relativistic limit, both unpolarized DCSs (non relativistic and relativistic) give very close results. We also gave the envelopes of the two unpolarized DCSs in both regimes for the sake of illustration and also to retrieve the shape of Fig. (3) of [11] which was confirmed since the scaling is the same (10^{-7}) and the asymmetric deviation from the elastic peak was also found as expected. The envelope for the relativistic regime was more difficult to obtain since the stability of the very robust code that evaluates the ordinary Bessel functions was put on trial because of the very high or very low arguments of the Bessel functions. Needless to say that working in simple precision, the results obtained are nonetheless sound and coherent.

VI. APPENDIX

$$\begin{aligned} \mathcal{A}(\lambda_i, \lambda_f) = & \left\{ 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * |p_f|^2 * |p_i|^2 * c^8 * \cos(\theta_{if}) - 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * |p_f|^2 * c^6 * \cos(\theta_{if}) * E_i^2 + \right. \\ & 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * |p_f| * |p_i| * c^{10} - 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * |p_i|^2 * c^6 * \cos(\theta_{if}) * E_f^2 + 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \\ & \lambda_i * c^8 * \cos(\theta_{if}) * E_f * E_i + 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * c^4 * \cos(\theta_{if}) * E_f^2 * E_i^2 + 2.0 * (k.p_f)^2 * (k.p_i)^2 * |p_f| * |p_i| * c^{10} * \cos(\theta_{if}) + \\ & 2.0 * (k.p_f)^2 * (k.p_i)^2 * c^{12} + 2.0 * (k.p_f)^2 * (k.p_i)^2 * c^8 * E_f * E_i + 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * (a^2) * |p_i|^2 * c^4 * \cos(\theta_{if}) * E_f * \omega - \\ & 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * (a^2) * c^2 * \cos(\theta_{if}) * E_f * E_i^2 * \omega - 2.0 * (k.p_f)^2 * (k.p_i)^2 * (a^2) * c^6 * E_i * \omega + 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \\ & \lambda_i * (a^2) * |p_f|^2 * c^4 * \cos(\theta_{if}) * E_i * \omega - 2.0 * (k.p_f)^2 * (k.p_i)^2 * \lambda_f * \lambda_i * (a^2) * c^2 * \cos(\theta_{if}) * E_f^2 * E_i * \omega - 2.0 * (k.p_f)^2 * (k.p_i)^2 * (a^2) * \end{aligned}$$

$$\begin{aligned}
& (a^2)*c^{10}+(k.p_f)^2*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*c^6*E_f*E_i-2.0*(k.p_f)^2*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|^2*|p_i|*c^6*\cos(\theta_{if})*\omega- \\
& 2.0*(k.p_f)^2*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^8*\omega+2.0*(k.p_f)^2*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^4*\cos(\theta_{if})*E_f^2*\omega-(k.p_f)^2*(k.s_i)*\lambda_i* \\
& (a^2)*|p_f|*c^8*\cos(\theta_f)*\omega+(k.p_f)^2*(k.s_i)*\lambda_i*(a^2)*|p_i|*c^8*\cos(\theta_i)*\omega-(k.p_f)^2*(k.s_i)*\lambda_i*(a^2)*c^7*E_f*\omega-(k.p_f)^2*(k.s_i)* \\
& \lambda_i*(a^2)*c^7*E_i*\omega+(k.p_f)*(k.p_i)^3*\lambda_f*\lambda_i*(a^2)*|p_f|*|p_i|*c^6+(k.p_f)*(k.p_i)^3*\lambda_f*\lambda_i*(a^2)*c^4*\cos(\theta_{if})*E_f*E_i+(k.p_f)* \\
& (k.p_i)^3*(a^2)*c^8+2.0*(k.p_f)*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^6*\cos(\theta_{if})*E_i+2.0*(k.p_f)*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)* \\
& |p_i|*c^6*E_f-(k.p_f)*(k.p_i)^2*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^6*E_i-(k.p_f)*(k.p_i)^2*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^6*\cos(\theta_{if})*E_f- \\
& 2.0*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*(a^2)*|p_f|^2*c^4*\cos(\theta_{if})*E_i*\omega+2.0*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*(a^2)*|p_i|^2*c^4*\cos(\theta_{if})*E_f*\omega+ \\
& 2.0*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*(a^2)*c^2*\cos(\theta_{if})*E_f^2*E_i*\omega-2.0*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*(a^2)*c^2*\cos(\theta_{if})*E_f*E_i^2*\omega+2.0* \\
& (k.p_f)*(k.p_i)^2*\lambda_f*(a^2)*|p_f|*c^7*\omega-2.0*(k.p_f)*(k.p_i)^2*\lambda_i*(a^2)*|p_i|*c^7*\omega+2.0*(k.p_f)*(k.p_i)^2*(a^2)*c^6*E_f*\omega-2.0* \\
& (k.p_f)*(k.p_i)^2*(a^2)*c^6*E_i*\omega-2.0*(k.p_f)*(k.p_i)*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|*|p_i|*c^8*\cos(\theta_{if})-2.0*(k.p_f)*(k.p_i)* \\
& (k.s_f)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*c^{10}-2.0*(k.p_f)*(k.p_i)*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*c^6*E_f*E_i+2.0*(k.p_f)*(k.p_i)*(k.s_f)*\lambda_f* \\
& \lambda_i*(a^2)*|p_f|*|p_i|^2*c^6*\cos(\theta_{if})*\omega-2.0*(k.p_f)*(k.p_i)*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^4*\cos(\theta_{if})*E_i^2*\omega+2.0*(k.p_f)*(k.p_i)* \\
& (k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^8*\omega+2.0*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|^2*|p_i|*c^6*\cos(\theta_{if})*\omega+2.0*(k.p_f)*(k.p_i)* \\
& (k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^8*\omega-2.0*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^4*\cos(\theta_{if})*E_f^2*\omega+2.0*(k.p_f)*(k.p_i)*(k.s_i)* \\
& \lambda_i*(a^2)*c^7*E_f*\omega+2.0*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_i*(a^2)*c^7*E_i*\omega-2.0*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*|p_f|^2*|p_i|^2*c^4*\cos(\theta_{if})* \\
& \omega^2+2.0*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*|p_f|^2*c^2*\cos(\theta_{if})*E_i^2*\omega^2-2.0*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*|p_f|*|p_i|*c^6*\omega^2+2.0* \\
& (k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*|p_i|^2*c^2*\cos(\theta_{if})*E_f^2*\omega^2+2.0*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*c^4*\cos(\theta_{if})*E_f*E_i*\omega^2-2.0* \\
& (k.p_f)*(k.p_i)*\lambda_f*\lambda_i*(a^2)*\cos(\theta_{if})*E_f^2*E_i*\omega^2-2.0*(k.p_f)*(k.p_i)*\lambda_f*(a^2)*|p_f|^2*c^6*\cos(\theta_f)*\omega^2+2.0*(k.p_f)*(k.p_i)* \\
& \lambda_f*(a^2)*|p_f|*|p_i|*c^6*\cos(\theta_i)*\omega^2-2.0*(k.p_f)*(k.p_i)*\lambda_f*(a^2)*|p_f|*c^5*E_i*\omega^2+2.0*(k.p_f)*(k.p_i)*\lambda_f*(a^2)*c^4*\cos(\theta_f)* \\
& E_f^2*\omega^2-2.0*(k.p_f)*(k.p_i)*\lambda_i*(a^2)*|p_i|*c^5*E_f*\omega^2+2.0*(k.p_f)*(k.p_i)*\lambda_i*(a^2)*c^4*\cos(\theta_i)*E_f*E_i*\omega^2+2.0*(k.p_f)* \\
& (k.p_i)*(a^2)*|p_f|*|p_i|*c^6*\cos(\theta_{if})*\omega^2+2.0*(k.p_f)*(k.p_i)*(a^2)*c^8*\omega^2-2.0*(k.p_f)*(k.p_i)*(a^2)*c^4*E_f*E_i*\omega^2-(k.p_i)^3* \\
& (k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^6*\cos(\theta_{if})*E_i-(k.p_i)^3*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^6*E_f-(k.p_i)^3*\lambda_f*(a^2)*|p_f|*c^7*\omega+(k.p_i)^3* \\
& \lambda_f*(a^2)*c^6*\cos(\theta_f)*E_f*\omega+(k.p_i)^3*\lambda_i*(a^2)*|p_i|*c^7*\omega+(k.p_i)^3*\lambda_i*(a^2)*c^6*\cos(\theta_i)*E_i*\omega+(k.p_i)^2*(k.s_f)*(k.s_i)*\lambda_f* \\
& \lambda_i*(a^2)*|p_f|*|p_i|*c^8*\cos(\theta_{if})+(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)*c^{10}+(k.p_i)^2*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*(a^2)*c^6*E_f*E_i- \\
& 2.0*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_f|*|p_i|^2*c^6*\cos(\theta_{if})*\omega+2.0*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_f|*c^4*\cos(\theta_{if})*E_i^2*\omega- \\
& 2.0*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*(a^2)*|p_i|*c^8*\omega+(k.p_i)^2*(k.s_i)*\lambda_i*(a^2)*|p_f|*c^8*\cos(\theta_f)*\omega-(k.p_i)^2*(k.s_i)*\lambda_i*(a^2)*|p_i|*c^8* \\
& \cos(\theta_i)*\omega-(k.p_i)^2*(k.s_i)*\lambda_i*(a^2)*c^7*E_f*\omega-(k.p_i)^2*(k.s_i)*\lambda_i*(a^2)*c^7*E_i*\omega+2.0*(k.p_i)^2*\lambda_f*(a^2)*|p_f|*c^5*E_i*\omega^2- \\
& 2.0*(k.p_i)^2*\lambda_f*(a^2)*c^4*\cos(\theta_f)*E_f*E_i*\omega^2-2.0*(k.p_i)^2*\lambda_i*(a^2)*|p_f|*|p_i|*c^6*\cos(\theta_f)*\omega^2+2.0*(k.p_i)^2*\lambda_i*(a^2)*|p_i|^2* \\
& c^6*\cos(\theta_i)*\omega^2+2.0*(k.p_i)^2*\lambda_i*(a^2)*|p_i|*c^5*E_f*\omega^2-2.0*(k.p_i)^2*\lambda_i*(a^2)*c^4*\cos(\theta_i)*E_i^2*\omega^2\}/[4.0*(k.p_f)^2*(k.p_i)^2*c^8]
\end{aligned}$$

$$\begin{aligned}
\mathcal{C}(\lambda_i, \lambda_f) = & \left\{ \cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)^2*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)^2* \right. \\
& (k.s_i)*\lambda_f*\lambda_i*|p_f|*c^4*\omega-2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)*(k.p_i)*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.p_f)* \\
& (a_1.p_i)*(k.p_f)*(k.p_i)*(k.s_f)*\lambda_f*\lambda_i*|p_i|*c^4*\omega+2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*|p_f|*c^4*\omega- \\
& 2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*|p_f|*|p_i|*c^2*\omega^2+2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)*(k.p_i)*\lambda_f* \\
& \lambda_i*\cos(\theta_{if})*E_f*E_i*\omega^2+2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_f)*(k.p_i)*c^4*\omega^2+\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_i)^2*(k.s_f)* \\
& (k.s_i)*\lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*|p_i|*c^4*\omega-\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_i)^2* \\
& (k.p_i)*(k.s_f)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.s_f)*(k.p_f)^2*(k.p_i)*\lambda_f*\lambda_i*|p_i|*c^2*\cos(\theta_{if})*E_f*\omega^2-\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_i)^3* \\
& (k.s_f)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.p_f)*(a_1.p_i)*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*c^4*E_i*\omega-\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)^3* \\
& (k.s_i)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)^2*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)^2* \\
& (k.p_i)*\lambda_f*\lambda_i*|p_i|*c^4*\omega+2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)^2*(k.s_i)*\lambda_f*\lambda_i*c^4*E_f*\omega-\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)* \\
& (k.p_f)*(k.p_i)^2*(k.s_i)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*|p_i|*c^4*\omega-2.0*\cos(2\phi_0)*(a_1.p_i)* \\
& (a_1.s_f)*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*c^4*E_f*\omega-2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*|p_f|*c^2*\cos(\theta_{if})* \\
& E_i*\omega^2+2.0*\cos(2\phi_0)*(a_1.p_i)*(a_1.s_f)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*|p_i|*c^2*E_f*\omega^2+\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)^3*(k.p_i)* \\
& \lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)^2*(k.p_i)^2*\lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)^2*(k.p_i)* \\
& \lambda_f*\lambda_i*c^4*E_f*\omega+2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)^2*(k.p_i)*\lambda_f*\lambda_i*c^4*E_i*\omega+\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)* \\
& (k.p_i)^3*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*c^4*E_f*\omega-2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)* \\
& (k.p_f)*(k.p_i)^2*\lambda_f*\lambda_i*c^4*E_i*\omega+2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*|p_f|*|p_i|*c^4*\cos(\theta_{if})*\omega^2+2.0* \\
& \cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*c^6*\omega^2-2.0*\cos(2\phi_0)*(a_1.s_f)*(a_1.s_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*c^2*E_f*E_i* \\
& \omega^2-\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)^2*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*c^6+2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)^2*(k.s_i)*\lambda_f*\lambda_i* \\
& |p_f|*c^4*\omega+2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)*(k.p_i)*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i*c^6-2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)* \\
& (k.p_i)*(k.s_f)*\lambda_f*\lambda_i*|p_i|*c^4*\omega-2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)*(k.p_i)*(k.s_i)*\lambda_f*\lambda_i*|p_f|*c^4*\omega+2.0*\cos(2\phi_0)* \\
& (a_2.p_f)*(a_2.p_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*|p_f|*|p_i|*c^2*\omega^2-2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)*(k.p_i)*\lambda_f*\lambda_i*\cos(\theta_{if})*E_f* \\
& E_i*\omega^2-2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)*(k.p_i)*c^4*\omega^2-\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_i)^2*(k.s_f)*(k.s_i)*\lambda_f*\lambda_i* \\
& c^6+2.0*\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_i)^2*(k.s_f)*\lambda_f*\lambda_i*|p_i|*c^4*\omega+\cos(2\phi_0)*(a_2.p_f)*(a_2.p_i)*(k.p_f)^2*(k.p_i)*(k.s_f)*
\end{aligned}$$

$$\begin{aligned}
& \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_f * c^5 * E_f^2 * \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_f * c^5 * E_f * \\
& E_i * \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_i * |p_f| * |p_i| * c^7 * \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \\
& \lambda_i * |p_f| * c^6 * \cos(\theta_f) * E_i * \omega \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_i * |p_f| * c^6 * \cos(\theta_i) * E_i * \omega \sin(\varphi_f) \sin(\theta_f) * |a| - \\
& \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_i * |p_i|^2 * c^7 * \omega \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_i * c^5 * E_f * E_i * \omega \sin(\varphi_i) \sin(\theta_i) * \\
& |a| + \sin(\phi_0) * (k.p_f)^2 * (k.p_i) * \lambda_i * c^5 * E_i^2 * \omega \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_f * |p_f|^2 * c^7 * \omega \sin(\varphi_f) \sin(\theta_f) * \\
& |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_f * |p_f| * |p_i| * c^7 * \omega \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_f * |p_i|^2 * c^6 * \cos(\theta_f) * E_f * \\
& \omega \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_f * |p_i|^2 * c^6 * \cos(\theta_i) * E_f * \omega \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \\
& \lambda_f * c^5 * E_f^2 * \omega \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_f * c^5 * E_f * E_i * \omega \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_f) * \\
& (k.p_i)^2 * \lambda_i * |p_f| * |p_i| * c^7 * \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_i * |p_f| * c^6 * \cos(\theta_f) * E_i * \omega \sin(\varphi_i) \sin(\theta_i) * \\
& |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_i * |p_f| * c^6 * \cos(\theta_i) * E_i * \omega \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_i * |p_i|^2 * c^7 * \omega * \\
& \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_i * c^5 * E_f * E_i * \omega \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i)^2 * \lambda_i * c^5 * E_i^2 * \\
& \omega \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i) * \lambda_f * (a^2) * c^3 * E_f * \omega^2 \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_f) * (k.p_i) * \lambda_i * (a^2) * \\
& c^3 * E_i * \omega^2 \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_f) * (k.s_i) * \lambda_i * (a^2) * |p_f| * c^5 * \omega^2 \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_f) * (k.s_i) * \\
& \lambda_i * (a^2) * |p_i| * c^5 * \omega^2 \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_i)^2 * \lambda_f * (a^2) * c^3 * E_f * \omega^2 \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_i)^2 * \lambda_i * \\
& (a^2) * c^3 * E_i * \omega^2 \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_i) * (k.s_i) * \lambda_i * (a^2) * |p_f| * c^5 * \omega^2 \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_i) * \\
& (k.s_i) * \lambda_i * (a^2) * |p_i| * c^5 * \omega^2 \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_f * (a^2) * |p_f|^2 * c^3 * \omega^3 \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * \\
& (k.p_i) * \lambda_f * (a^2) * |p_f| * |p_i| * c^3 * \omega^3 \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_f * (a^2) * |p_i| * c^2 * \cos(\theta_f) * E_f * \omega^3 \sin(\varphi_i) \sin(\theta_i) * \\
& |a| + \sin(\phi_0) * (k.p_i) * \lambda_f * (a^2) * |p_i| * c^2 * \cos(\theta_i) * E_f * \omega^3 \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_i) * \lambda_f * (a^2) * c * E_f^2 * \omega^3 \sin(\varphi_f) * \\
& \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_f * (a^2) * c * E_f * E_i * \omega^3 \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_i * (a^2) * |p_f| * |p_i| * c^3 * \omega^3 * \\
& \sin(\varphi_f) \sin(\theta_f) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_i * (a^2) * |p_f| * c^2 * \cos(\theta_f) * E_i * \omega^3 \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_i) * \lambda_i * (a^2) * |p_f| * \\
& c^2 * \cos(\theta_i) * E_i * \omega^3 \sin(\varphi_f) \sin(\theta_f) * |a| + \sin(\phi_0) * (k.p_i) * \lambda_i * (a^2) * |p_i|^2 * c^3 * \omega^3 \sin(\varphi_i) \sin(\theta_i) * |a| + \sin(\phi_0) * (k.p_i) * \lambda_i * \\
& (a^2) * c * E_f * E_i * \omega^3 \sin(\varphi_i) \sin(\theta_i) * |a| - \sin(\phi_0) * (k.p_i) * \lambda_i * (a^2) * c * E_i^2 * \omega^3 \sin(\varphi_i) \sin(\theta_i) * |a| \} / [2.0 * (k.p_f)^2 * (k.p_i)^2 * c^7]
\end{aligned}$$

-
- [1] T. H. Maiman, *Nature*, **187**, 493-494, (1961).
[2] M. H. Mittleman, *Introduction to The Theory of Laser-Atom Interactions*, Springer, 2ed, (1993).
[3] Landau and E. M Lifshitz, *Quantum Mechanics, Non Relativistic Theory*, 3ed, New York Pergamon, (1997).
[4] W. Greiner, *Relativistic Quantum Mechanics*, 3ed, Springer, (2000); W. Greiner and J. Reinhardt *Quantum Electrodynamics*, 3ed, Springer, (2003).
[5] R. P. Feynman, R. B. Leighton and M. Sands, *The Feynman Lectures on Physics*, Volume **3**, Addison-Wesley, (1965).
[6] D. T. Pierce *et al*, *Rev. Sci. Instrum.*, **51**, 478, (1980); D. T. Pierce *et al*, *Appl. Phys. Lett.*, **26**, 670, (1975).
[7] J. Kessler, *Polarized Electrons*, 2ed, Springer, (1985).
[8] A. G. Grozin, *Using Reduce in High Energy Physics*, Cambridge University Press, Cambridge, 1997.
[9] B. Manaut *et al*, *Phys. Rev. A* **71**, 043401 (2005).
[10] F. V. Bunkin and M. V. Fedorov, *Sov. Phys. JETP* **22**, 844, 1966; N. M. Kroll and K. N. Watson, *Phys. Rev. A*, **8**, 804, 1973.
[11] C. Szymannowski *et al*, *Phys. Rev. A*, **56**, 3846, 1997.